

Title: Morphological Features of Pistol Grip Deformity

Authors

Kunihide MURAOKA^a, Masatoshi NAITO^a, Koichi KINOSHITA^a, Yoshinari
NAKAMURA^b, Tomonobu HAGIO^a

Affiliation

^aDepartment of Orthopaedic Surgery, Fukuoka University Faculty of Medicine

^bDepartment of Orthopaedic Surgery, Iida Hospital

Running Title: Morphological Features of PGD

Abstract

Purpose The purpose of this study was to elucidate the morphological features of the pistol grip deformity (PGD) using radial computed tomography (CT) images.

Materials and Methods Using multiplanar reformation, radial CT images of a total of 51 hips were reconstructed. The femoral neck was used as the axis of rotation and the images were generated at 10° intervals from anterior to posterior. The alpha angles of each plane were measured, and compared between the PGD and control (normal morphology) groups. We also compared the mean alpha angles between each plane within the PGD group.

Results The mean maximum alpha angles were 61° in the 60° vector in the PGD group and 50° in the 40° and 50° vectors in the control group. The mean alpha angles in the PGD group was significantly greater than those in the control group in the 30° to 100° vectors ($P<0.05$). In the PGD group, the mean alpha angles at the anterosuperior position were significantly greater than those at the superior position ($P<0.05$).

Discussion In this study, we found that PGD is a particular morphological abnormality from the anterosuperior to superior femoral head-neck junction. This is different from pure cam-type deformities that have previously been reported. We need to differentiate between these two types of morphological features, PGD and pure cam-type deformity, to understand the morphology for preoperative and intraoperative decision-making for femoroacetabular impingement.

Keywords Femoroacetabular impingement, Cam-type deformity, Pistol grip deformity, Radial imaging

Footnotes

Correspondence to: Kunihide MURAOKA

Department of Orthopaedic Surgery, Fukuoka University Faculty of Medicine, 7-45-1

Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan

Tel: +81-92-801-1011

Fax: +81-92-864-9055

E-mail: kagoshimaizn@yahoo.co.jp

Introduction

Femoroacetabular impingement (FAI) is the most common mechanism underlying the development of early cartilage and labral damage in the nondysplastic hip ¹⁾⁻³⁾. The most common femoral structural deformities involve loss of the femoral head-neck offset (cam-type deformities). Because there is evidence that incomplete correction of the femoral head-neck offset remains a leading cause of revision surgery for FAI ⁴⁾, a detailed preoperative understanding of the morphology is important for preoperative decision-making ⁵⁾.

A cam-type deformity can usually be detected on lateral radiographs of the hip. However, there is also a lesion that can be detected on anteroposterior (AP) radiographs of the hip, the so-called pistol grip deformity (PGD) ⁶⁾. The two types of morphological features (PGD and cam-type deformity) have been assessed in the same cohorts to date (Fig. 1, 2). However, we believe that they should not be confused, because they have obviously different morphological features on AP radiographs.

The aim of this study was to elucidate the morphological features of PGD using radial computed tomography (CT) images. To our knowledge, this is the first study to focus on the morphological features of PGD.

Materials and Methods

Fifty-one consecutive hips in 48 patients (31 men, 16 women; mean age, 63 years; age range, 30–96 years) who met our inclusion criteria were included after a retrospective review of the radiology information system and patient records. The inclusion criteria for the two groups were as follows. The PGD group was defined as obvious presence of loss of femoral head-neck offset on AP radiographs, triangular index ⁷⁾ of ≥ 2 mm, and

Tönnis grade ⁸⁾ of ≤ 1 . The control group (normal morphology) was defined as no history of ongoing hip or groin pain, no prior hip surgery, obvious absence of loss of femoral head-neck offset on AP radiographs, triangular index of < 2 mm, lateral center-edge angle of $\geq 25^\circ$, alpha angle ⁹⁾ on lateral radiographs of $< 50.5^\circ$, and Tönnis grade of ≤ 1 . Patients who had not undergone CT imaging of the hip joint were excluded even if they met our inclusion criteria.

A supine AP digital radiograph of the pelvis was obtained. The tube-to-film distance was 120 cm, with the tube perpendicular to the table. The center beam was directed toward the midpoint between the upper border of the symphysis and a horizontal line connecting the bilateral AP iliac spines. The radiographic measurements were performed using a software program (Rapideye™ Hyper; Toshiba, Tochigi, Japan). The CT images were obtained with an Aquilion ONE ViSION Edition (Toshiba, Ootawara, Japan), and multiplanar reformation was carried out with a ZIOSTATION (Ziosoft, Tokyo, Japan). Radial CT images of each hip were reconstructed and generated at 10° intervals from anterior to posterior (0° to 180°) of the femoral head, using the femoral neck as the axis of rotation (Figs. 3, 4). The alpha angles of each plane were measured. A value of $> 50.5^\circ$ was considered positive for cam-type morphology according to previous reports ⁹⁾⁻¹⁴⁾. We defined the 50° and 60° vectors as anterosuperior positions and the 80° and 90° vectors as superior positions based on previous studies ^{10), 15)-17)}.

For each group, the vector at which the mean maximum alpha angle measurement occurred and the percentage distribution of positive cam-type deformities at each plane were noted. The differences in the mean alpha angle values between the PGD group and the control group at each plane were determined using a two-sample *t*-test or Mann–Whitney U test depending on the normality of the distribution of the alpha angles

evaluated by the Shapiro–Wilk test. Analysis of variance was used to evaluate the differences in the mean alpha angle values at each plane within the PGD group. Finally, the rates of positive cam-type morphology at each plane were compared between the two groups using the chi-square test. Values of $P < 0.05$ were considered statistically significant.

The alpha angle measurements were performed by two of the authors, and the intraclass correlation coefficients (ICCs) were determined.

Results

In the PGD group, there were 31 hips in 8 female and 20 male patients with a mean age of 65 years. In the control group, there were 20 hips in 8 female and 12 male patients with a mean age of 60 years. The mean maximum alpha angles were 61° in the 60° vector in the PGD group and 50° in the 40° and 50° vectors in the control group. The mean alpha angles in the PGD group were significantly greater than those in the control group in the 30° to 100° vectors (Table 1). In the PGD group, the mean alpha angles in the 50° and 60° vectors (anterosuperior positions) were significantly greater than those in the 90° to 100° vectors and 80° to 100° vectors (superior positions), respectively. In the PGD group, the rates of positive cam-type morphology were significantly higher than those in the normal group in the 30° to 90° vectors (Table 2).

The ICCs for measuring the alpha angles were 0.992 (95% confidence interval [CI], 0.991 to 0.993) for intraobserver reliability and 0.969 (95% CI, 0.964 to 0.972) for interobserver reliability.

Discussion

In cam-type deformities, there are two obviously different types of morphological abnormalities. These comprise abnormalities that can even be seen on AP radiographs (PGD) and only seen on lateral radiographs (pure cam-type deformity). Although PGD was originally described by Stulberg et al. ⁶⁾ as a morphological feature that can be seen on AP radiographs, Eijer et al. ¹⁸⁾ subsequently reported that it can be best identified on lateral radiographs. Since then, most studies evaluating cam-type deformities have assessed these two types of deformities in confusion. Because there is evidence that incomplete correction of the femoral head-neck offset remains the leading cause of revision surgery for FAI ⁴⁾, a detailed preoperative understanding of the morphology is important for preoperative decision-making ⁵⁾. We believe that these two types of morphological features (PGD and pure cam-type deformity) should not be confused because they have obviously different morphological features on AP radiographs.

There are some studies that have addressed the morphological features of cam-type deformities using radial images. Siebenrock et al. ¹⁷⁾ described that the FAI group showed a statistically significant decrease in the femoral head-neck offset in the anterosuperior quadrant compared with the control group. Phirmann et al. ¹⁶⁾ reported that the alpha angle was significantly larger in patients with cam-type FAI at the anterior and anterosuperior positions. To the best of our knowledge, no studies have reported the morphological abnormalities in the superior femoral head-neck junction in FAI hips. In this study, we found that the mean alpha angles in the PGD group were significantly greater than those in the control group in the 30° to 100° vectors and that the rates of positive cam-type deformities were significantly higher in the PGD group than in the normal group in the 30° to 90° vectors. The PGD group was revealed to have cam-type deformities not only at the anterosuperior but also at the superior femoral head-neck

junction. In FAI hips, Tamura et al.¹⁹⁾ described that a mode 1 labral tear, which may be a risk factor for the development of adjacent acetabular cartilage damage, was likely to occur at the anterior and anterosuperior zones. Our findings imply that the cartilage and labral damage in PGD hips may occur in a wider range of zones compared with hips with pure cam-type deformities, and that a wider range of resection of the femoral head-neck junction may be needed for PGD hips. Further investigations to evaluate the differences in the location and degree of cartilage or labral damage between hips with PGD and pure cam-type deformity will be needed.

In the PGD group, the mean maximum alpha angle was 61° in the 60° vector. This finding was similar to previous reports that have been published^{15), 17)}. In the PGD group, we also found that the mean alpha angles at the anterosuperior position were significantly greater than those at the superior position. The percentage distributions of positive cam-type morphology at the 50°, 60°, 80°, 90°, and 100° vectors were 90%, 90%, 61%, 39%, and 10%, respectively. Therefore, PGD hips had a high rate of cam-type deformities at anterosuperior positions. Meanwhile, they also had cam-type deformities at superior positions at lower rates. Rakhra et al.¹⁵⁾ reported that radial plane imaging should be considered to avoid underestimating or missing the morphological abnormalities of FAI hips. Our study supports their opinion. Because there is a possibility of overlooking cam-type deformities, especially at the superior position, in PGD hips without three-dimensional evaluations, we recommend radial imaging evaluation for preoperative decision-making for cam-type FAI.

There is a limitation to our study. The alpha angles of pure cam-type deformity should ideally be compared with those of PGD. However, it was difficult for us to collect a sufficiently large sample size for pure cam-type FAI because of the low incidence of

pure FAI in Japan²⁰⁾. However, we feel that our study design is not inferior to those in previous reports evaluating the morphological features of cam-type FAI^{15), 16)}.

In conclusion, we found that PGD is a particular morphological abnormality from the anterosuperior to superior femoral head-neck junction, which is different from pure cam-type deformities that have previously been described. For preoperative and intraoperative decision-making for cam-type FAI we need to differentiate these two types of morphological features, and a radial imaging evaluation is recommended.

References

1. Beck M, Kalhor M, Leunig M, Ganz R. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br* 87: 1012-1018, 2005
2. Clohisy JC, St John LC, Schutz AL. Surgical treatment of femoroacetabular impingement: a systematic review of the literature. *Clin Orthop Relat Res* 468: 555-564, 2010
3. Tanzer M, Noiseux N. Osseous abnormalities and early osteoarthritis: the role of hip impingement. *Clin Orthop Relat Res* 429: 170-177, 2004
4. Philippon MJ, Schenker ML, Briggs KK, Kuppersmith DA, Maxwell RB, Stubbs AJ. Revision hip arthroscopy. *Am J Sports Med* 35: 1918-1921, 2007
5. Ipach I, Arlt EM, Mittag F, Kunze B, Wolf P, Kluba T. A classification-system improves the intra- and interobserver reliability of radiographic diagnosis of "pistol-grip-deformity". *Hip Int* 21: 732-739, 2011
6. Stulberg S, Cordell L, Harris W, Ramsey P, MacEwen G. Unrecognized Childhood Hip Disease: A Major Cause of Idiopathic Osteoarthritis of the Hip. In: Society TH, ed. *Third Open Scientific Meeting of The Hip Society*. Saint Louis: CV Mosby. 212. 1975
7. Gosvig KK, Jacobsen S, Palm H, Sonne-Holm S, Magnusson E. A new radiological index for assessing asphericity of the femoral head in cam impingement. *J Bone Joint Surg Br* 89: 1309-1316, 2007
8. Tönnis D, Heinecke A. Acetabular and femoral anteversion: relationship with osteoarthritis of the hip. *J Bone Joint Surg Am* 81: 1747-1770, 1999
9. Nötzli HP, Wyss TF, Stoecklin CH, Schmid MR, Treiber K, Hodler J. The contour of the femoral head-neck junction as a predictor for the risk of anterior impingement. *J Bone Joint Surg Br* 84: 556-560, 2002

10. Hack K, Di Primio G, Rakhra K, Beaulé PE. Prevalence of cam-type femoroacetabular impingement morphology in asymptomatic volunteers. *J Bone Joint Surg Am* 92: 2436-2444, 2010
11. Meyer DC, Beck M, Ellis T, Ganz R, Leunig M. Comparison of six radiographic projections to assess femoral head/neck asphericity. *Clin Orthop Relat Res* 445: 181-185, 2006
12. Tannast M, Siebenrock KA, Anderson SE. Femoroacetabular impingement: radiographic diagnosis--what the radiologist should know. *AJR Am J Roentgenol* 188: 1540-1552, 2007
13. Beaulé PE, Zaragoza E, Motamedi K, Copelan N, Dorey FJ. Three-dimensional computed tomography of the hip in the assessment of femoroacetabular impingement. *J Orthop Res* 23: 1286-1292, 2005
14. Barton C, Salineros MJ, Rakhra KS, Beaulé PE. Validity of the alpha angle measurement on plain radiographs in the evaluation of cam-type femoroacetabular impingement. *Clin Orthop Relat Res* 469: 464-469, 2011
15. Rakhra KS, Sheikh AM, Allen D, Beaulé PE. Comparison of MRI alpha angle measurement planes in femoroacetabular impingement. *Clin Orthop Relat Res* 467: 660-665, 2009
16. Pfirrmann CW, Mengiardi B, Dora C, Kalberer F, Zanetti M, Hodler J. Cam and pincer femoroacetabular impingement: characteristic MR arthrographic findings in 50 patients. *Radiology* 240: 778-785, 2006
17. Siebenrock KA, Wahab KH, Werlen S, Kalhor M, Leunig M, Ganz R. Abnormal extension of the femoral head epiphysis as a cause of cam impingement. *Clin Orthop Relat Res* 418: 54-60, 2004
18. Eijer H, Leunig M, Mahomed N, Ganz R. Cross-table lateral radiographs for screening of anterior femoral head-neck offset in patients with femoroacetabular impingement. *Hip Int* 11: 37-41, 2001
19. Tamura S, Nishii T, Takao M, Sakai T, Yoshikawa H, Sugano N. Differences in the locations and modes of labral tearing between dysplastic hips and those with femoroacetabular impingement. *Bone Joint J* 95-B: 1320-1325, 2013
20. Takeyama A, Naito M, Shiramizu K, Kiyama T. Prevalence of femoroacetabular impingement in Asian patients with osteoarthritis of the hip. *Int Orthop* 33:1229-1232, 2009

Legends for Figures

Fig. 1 AP and lateral radiographs of hips showing a typical case of PGD

Fig. 2 AP and lateral radiographs of hips showing a typical case of pure cam-type deformity

Fig. 3 The panel shows the process for reconstructing radial images from anterior (0°) to posterior (180°) of the femoral head at 10° intervals, using the femoral neck as the axis of rotation

Fig. 4 Radial images at each vector

Figures




Fig. 1 AP and lateral radiographs of hips showing a typical case of PGD



Fig. 2 AP and lateral radiographs of hips showing a typical case of pure cam-type deformity




Fig. 3 The panel shows the process for reconstructing radial images from anterior (0°) to posterior (180°) of the femoral head at 10° intervals, using the femoral neck as the axis of rotation




Fig. 4 Radial images at each vector

Tables

Table 1 Comparison of the alpha angles of each plane between the two groups

| Plane (°) | | 0 | 10 | 20 | 30 | 40 |
|---------------|--------------------|-----------|-----------|-----------|-----------|-----------|
| PGD group | Alpha angle (°) | | | | | |
| | mean | 40.5 | 43.6 | 47.3 | 52.7 | 57.8 |
| | range | 23.9-57.5 | 31.3-59.8 | 31.1-63.9 | 32.6-69.7 | 39.3-82.8 |
| | standard deviation | 8.1 | 8.1 | 8.4 | 9.3 | 9.9 |
| Control group | Alpha angle (°) | | | | | |
| | mean | 39.1 | 41.4 | 44.8 | 47.2 | 50.1 |
| | range | 32.2-47.3 | 33.8-49.8 | 35.3-54.9 | 37.8-57.7 | 39.7-60.0 |
| | standard deviation | 4.5 | 4.9 | 5.4 | 5.3 | 6.2 |
| P value | | | n.s. | | 0.011 | 0.001 |

| Anterosuperior | | | Superior | | | | |
|----------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 50 | 60 | 70 | 80 | 90 | 100 | 110 | 120 |
| 59.6 | 60.6 | 56.2 | 53.3 | 48.3 | 42.9 | 40.1 | 37.5 |
| 39.0-81.7 | 42.5-88.0 | 42.0-72.9 | 36.9-75.4 | 35.1-70.3 | 33.2-63.5 | 32.1-64.6 | 29.2-63.8 |
| 9.0 | 9.4 | 8.6 | 9.1 | 8.8 | 7.2 | 6.5 | 6.4 |
| 49.7 | 46.2 | 44.3 | 41.2 | 39.6 | 38 | 37.7 | 36.2 |
| 42.7-60.3 | 39.3-53.3 | 39.0-51.7 | 35.9-45.1 | 37.0-44.8 | 33.3-43.8 | 32.1-49.7 | 30.4-49.7 |
| 5.7 | 4.0 | 3.6 | 2.5 | 2.3 | 3.2 | 4.5 | 4.7 |
| <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | 0.005 | n.s. | |

| 130 | 140 | 150 | 160 | 170 | 180 |
|-----------|-----------|-----------|-----------|-----------|-----------|
| 35.8 | 34.1 | 33.5 | 32.3 | 31.5 | 31.3 |
| 27.5-46.4 | 27.7-42.9 | 26.1-50.8 | 23.8-45.3 | 22.1-39.8 | 19.9-41.5 |
| 4.5 | 3.4 | 4.4 | 4.4 | 4.6 | 5.5 |
| 34.3 | 34.1 | 33.4 | 33.2 | 32.4 | 31.6 |
| 28.7-48.1 | 28.9-41.1 | 26.8-40.1 | 26.6-46.6 | 27.7-48.6 | 26.4-47.4 |
| 5.1 | 3.4 | 3.3 | 4.4 | 4.6 | 4.7 |
| n.s. | | | | | |

n.s. = not significant.

Table 2 Percentage distribution of cam-type morphology at each plane and the differences between the two groups

| Plane (°) | No. of hips | | P value |
|-----------|-------------|---------------|---------|
| | PGD group | Control group | |
| 0 | 3 (10%) | 0 (0%) | n.s. |
| 10 | 6 (19%) | 0 (0%) | n.s. |
| 20 | 12 (39%) | 2 (11%) | n.s. |
| 30 | 21 (68%) | 4 (22%) | 0.002 |
| 40 | 24 (77%) | 9 (50%) | 0.048 |
| 50 | 28 (90%) | 8 (44%) | 0.001 |
| 60 | 28 (90%) | 4 (22%) | <0.001 |
| 70 | 21 (68%) | 1 (6%) | <0.001 |
| 80 | 19 (61%) | 0 (0%) | <0.001 |
| 90 | 12 (39%) | 0 (0%) | 0.002 |
| 100 | 3 (10%) | 0 (0%) | n.s. |

n.s. = not significant.