

Δ^9 -Tetrahydrocannabinol Treatment Enhances the Immobility Time during the Forced Swimming Test in Mice

Shozo CHIDORI¹, Nobuaki EGASIRA², Tomomi MATSUDA²,
Emi KOUSHI², Hiroshi NAGAI¹, Michiko MATSUSHITA¹,
Naoki UCHIDA¹, Kenichi MISHIMA², Katsunori IWASAKI²,
Michihiro FUJIWARA² and Ryoji NISHIMURA¹

¹) *Department of Psychiatry, Fukuoka University School of Medicine, Fukuoka University*

²) *Department of Neuropharmacology, Faculty of Pharmaceutical Sciences, Fukuoka University*

Abstract : In the present study, we examined the effect of Δ^9 -tetrahydrocannabinol (THC), the principal psychoactive component of marijuana, on the immobility time during the forced swimming test (FST), a test with a high predictivity of antidepressant efficacy in human depression, in mice. THC enhanced the immobility time in the FST at doses of 0.3, 1, 3, and 6 mg/kg (i.p.). On the other hand, cannabinol (CBN) and cannabidiol (CBD), the major nonpsychoactive components of marijuana, had no effect on the immobility time in the FST. Moreover, we also found that THC (6 mg/kg i.p.) had no effect on the motor function such as the locomotor activity or motor coordination in both open-field and rota-rod tests. SR141716, a CB₁ receptor antagonist, reversed the THC (6 mg/kg i.p.) induced enhancement of the immobility time in the FST at a dose of 3 mg/kg (i.p.). These findings therefore suggest that THC enhances the immobility time in the FST through CB₁ receptor.

Key words : Forced Swimming Test, Δ^9 -tetrahydrocannabinol, CB₁ receptor, SR141716