

Evaluation of Changes in Anxiety, Depression, Social Behavior, and Oxytocin mRNA Levels in Adults after Adolescent Interventions in Maternal-separated Mice

Reo HAYASHI¹⁾, Leo GOTOH^{1),2)}, Hikaru HORI¹⁾
Hitoshi IIDA¹⁾, Hiroaki KAWASAKI¹⁾

¹⁾ *Department of Psychiatry, Faculty of Medicine, Fukuoka University*

²⁾ *Laboratory of Neuroscience, Department of Psychiatry, Faculty of Medicine, Fukuoka University*

Abstract

It has been reported that the oxytocin-mediated signal transduction system is biologically altered by traumatic childhood experiences. Oxytocin has also been shown to be involved in stress, social behavior and attachment with others. The effects of oxytocin administration on abused children according to their developmental stage and the changes in the oxytocin signaling pathway due to oxytocin administration or attachment formation are largely unknown. In this study, we used maternal-separated (postnatal day [PD] 2-15) mice as a model for childhood adversity and investigated the effects of intervention during adolescence (PD 33-47) by continuously administering oxytocin (1mg/kg/day) or cohabitating them with littermates who had not experienced maternal separation. Cohabitation group was created to develop healthy relationship and attachment with other individuals. The effects of interventions on anxiety- and depression-associated behavior and social behavior in adult mice (N=57) were assessed using behavioral assays from PD60 to PD65. Oxytocin signaling was evaluated by quantification of *oxytocin* (*OXT*) and *oxytocin receptor* (*OXTR*) mRNA in the hypothalamus using RT-PCR. Results of our experimental separation and cohabitation protocols showed that the maternal-separated mice exhibited improved social behavior in social interaction test through each intervention. Biological analysis showed no significant changes in *OXT* and *OXTR* mRNA expression in the hypothalamus. Obtained result showed that oxytocin administration and cohabitation with healthy littermates to maternal-separated mice in adolescent improved social behavior in adulthood. However, the result of *OXT* and *OXTR* mRNA expression in the hypothalamus did not reveal significant changes through each intervention.

Key words: Oxytocin, Adolescence, Social behavior, Oxytocin mRNA

Introduction

Adversity experiences in early childhood are known to be a risk for a variety of mental illnesses, and many studies have shown an association with post-adult anxiety disorders, depression, substance abuse disorders, and personality disorders¹⁻³⁾. Reports also suggest that incompetent child care is associated with attachment anxiety and attachment avoidance in adult⁴⁾. Alterations

in oxytocin-mediated signaling systems have been reported as a biological result of childhood trauma⁵⁾. Oxytocin is a hormone synthesized in the hypothalamus and is widely known for its effects on uterine contractions and ejaculation. In the medical field, it has been used as a drug to induce labor and delivery. Recently, its involvement in social behavior, such as parent-child attachment behavior, has attracted attention⁶⁾. In healthy rats, oxytocin administered continuously before and after weaning and during adolescence has been reported to

result in improved social behavior, decreased anxious behavior, and increased *oxytocin receptor* mRNA levels in the hypothalamus after adulthood⁷⁾. Maternal separation is a frequently used model for childhood adversity in rodents⁸⁾. Although there is a report⁹⁾ that pre- and post-weaning administration of oxytocin to rats with maternal separation improved social behavior and, reduced anxious behavior, changes in the oxytocin system across different brain regions and changes in behavior depending on the timing of administration are largely unclear. In addition, when considering its application to actual clinical practice, it is thought to be difficult to administer oxytocin continuously on an outpatient basis to abused young children. In addition, the oxytocin system has been reported to be highly plasticized during the early postnatal period and adolescence¹⁰⁾. Thus, a clear need for understanding the effects and mechanism of action of oxytocin administration during adolescence. On the other hand, as an environmental intervention, it has been reported that children with attachment disorders due to child abuse reduced interpersonal problems and attachment avoidance and improved social behavior when they form attachments with others in a safe environment¹¹⁾. We assumed that oxytocin is involved in the mechanism of this improvement, however, changes in oxytocin signaling pathways in various brain regions are unknown. In this study, cohabitation was done to develop healthy relationship and attachment with other individuals. We continuously administered oxytocin or cohabitated maternal-separated mice with littermates during adolescence. Thus, we investigated the effects of interventions, on anxiety, depression and social behaviors as well as the changes in biological mechanisms.

Material & Methods

1. Animals

Pregnant C57BL/6J female mice were purchased from Charles River Laboratories Japan, Inc. Among these mice, 11 pregnant female mice were used for the experiment. One week after the pregnant mice arrived, they gave birth to 57 pups, which were used in the experiment. Food and water were available ad libitum. The vivarium was maintained at $21\pm 1^\circ\text{C}$. Room lights were switched to light/dark every 12 h (lit at 8:00 am. The research protocol was approved by the Fukuoka University Ethical Review Board on Animal Testing (approval number: 2106022).

2. Experimental design

The procedure for separation of mothers and pups was performed according to previous studies¹²⁾. They were divided into two groups based on their childhood environments. Twenty of the 57 animals (nine females and 11 males) were raised as the standard nest (SN), and 37 of the 57 (nineteen females and 18 males) were raised as the maternal separate (MS). In MS, mothers and infants were separated for 4 h per day from postnatal day (PD) 2 to PD5, and for 8 h per day from PD5 to PD16. When separating mothers from their pups, to reduce the psychological burden on the mothers, only the mothers were moved to another gauge first; then, the pups were moved to the gauge for separation, and the mothers were returned to their original gauge. To increase the stress load of maternal separation, early weaning was implemented from PD17 in MS, whereas in SN, normal weaning was implemented at PD21. In SN, the mother and pups were separated at PD21. In the MS, after early weaning, pups were transferred to individual gauges for rearing. Twelve of the 37 mice cohabitated with their siblings growing in the SN. Subsequently, the adolescent PD35 to PD48 pups underwent two interventions: continuous intraperitoneal administration of saline and intraperitoneal administration of oxytocin. Five groups standard nest-normal saline (SN-NS), maternal separate-normal saline (MS-NS), standard nest-oxytocin (SN-OXT), maternal separate-oxytocin (MS-OXT) and maternal separate-cohabitation with healthy littermates (MS-COHAB) were created according to childhood environment and type of injection or cohabitation. Behavioral analysis was conducted after PD60. After behavioral analysis, the mice were promptly decapitated, and their brains were removed and cryopreserved. Brain sections were then cut, RNA was extracted from the hypothalamus, and RT-PCR was performed. The schedule of the experiment is shown in Figure 1.

3. Drugs

Oxytocin was purchased from MERCK (Darmstadt, Germany). Oxytocin was dissolved in saline and administered intraperitoneally to the mice at a concentration of 1 mg/kg. Oxytocin concentration was determined based on a previous experiment⁷⁾.

4. Behavioral Assays

Open Field Test (OFT)

The OFT was used to measure general activity. This




Figure 1. Experimental design

Abbreviations: postnatal day (PD), standard nest (SN), maternal separate (MS), normal saline (NS), oxytocin (OXT)

procedure was based on a previous experiment¹³. The mouse was placed in the center of the apparatus (30 cm × 30 cm × 40 cm) and allowed to move freely for 5 min. The total distance traveled in the center was analyzed using the Panlab Smart ver 3.0.06.

Elevated Plus Maze Test (EPM)

The EPM was used to measure anxiety-associated behaviors. The procedure was based on a previous experiment¹⁴. The device consisted of an open arm (50 × 10 cm), a closed arm with a wall (50 × 10 × 40 cm), and a central square (10 × 10 cm) at a height of 60 cm. The time spent in the open arm was analyzed using the Panlab Smart ver 3.0.06.

Tail Suspension Test (TST)

The TST was used to measure depression-associated behavior. This procedure was based on a previous experiment¹⁵. The tail of the mouse was fixed to a rod installed in the test device so the mouse was hung upside-down. The immobility time was measured and analyzed using Panlab Smart ver 3.0.06.

Social Interaction Test (SIT)

The SIT was used to measure social behavior in terms of social proximity in mice. This procedure was based on a previous experiment¹⁶. First, the mice were placed in an open field (30 × 30 × 40 cm) with empty social interaction test cages and allowed to explore freely for 5 min. For the next 5 min, mice were allowed to explore freely in an open field (30 × 30 × 40 cm) containing social interaction test cages, including an unfamiliar mouse. The area close to the social interaction test cages was designated as high interaction zone and the area away from the cages as low interaction zone. The movements of the mice during the

first and the next 5 min were compared and analyzed using the Panlab Smart ver 3.0.06.

5. Biological analysis

RNA extraction and Real-time PCR

After behavioral testing, brain samples were removed and immediately frozen at -80 degrees and the hypothalamus was extracted from frozen brains at temperatures below freezing for RNA extraction. We determined to measure OXT and OXTR mRNA levels in the hypothalamus. The brain areas where the oxytocin system is evaluated generally include hypothalamus, nucleus accumbens (NAc), amygdala (AMY), hippocampus (Hip), prefrontal cortex (PFC), etc and the hypothalamus was selected in this study because it is the area where changes in oxytocin activity and *OXTR* mRNA were observed after MS or oxytocin administration on previous literature^{7,17}. Total RNA, excluding genomic DNA (gDNA), was extracted from separated brain samples using NucleoSpin[®] RNA (MACHEREY-NAGEL, Duren, Germany). Total RNA was reverse-transcribed to cDNA using SuperScript IV reverse transcriptase (Thermo Fisher Scientific, MA, USA). All samples were then amplified using THUNDERBIRD[®] SYBR[®] qPCR Mix (TOYOBO, Osaka, Japan) in a reaction volume of 20 μl. Referring to a previous study^{18,19}, *oxytocin* (*OXT*) and *oxytocin receptor* (*OXTR*) DNA were selectively amplified using each primer, and the *OXT* and *OXTR* DNA expression levels in the hypothalamus of each group were determined using *Actin beta* (*ACTB*) as an intrinsic control. All reactions were performed according to the manufacturer's protocol.

6. Statistical analysis

Analysis of variance (ANOVA) and Tukey-Kramer's honest significant difference (HSD) tests were used to compare the groups. Statistical significance was set at $P < 0.05$. All statistical analyses were performed using JMP 12.2.0 (SAS Institute Inc. Cary, NC).

Results

Open Field Test (OFT)

Analysis of variance for the total distance traveled showed a significant difference ($P < 0.01$). MS-OXT showed a significant increase compared to MS-COHAB ($P < 0.01$). MS-COHAB showed a significant decrease compared to MS-NS ($P < 0.01$). (Figure 2A)

SN-OXT showed no significant differences compared to SN-NS ($P = 0.1813$). (Figure 2B)




Figure 2. OFT Total distance traveled

Results from OFT. Total distance traveled in 5 minutes. Data are presented as means \pm SEM.

*: $p < 0.05$

Abbreviations: standard nest-normal saline (SN-NS), maternal separate-normal saline (MS-NS), maternal separate-oxytocin (MS-OXT), maternal separate-cohabitation with healthy littermates (MS-COHAB), standard nest-oxytocin (SN-OXT)

Elevated plus maze test (EPM)

Analysis of variance for the rate of time spent in the open arms showed no significant difference ($P = 0.26$). (Figure 3)




Figure 3. EPM Rate of time spent in the open arms

Results from EPM. Rate of time spent in the open arms in 5 minutes.

Data are presented as means \pm SEM.

*: $p < 0.05$

Abbreviations: standard nest-normal saline (SN-NS), maternal separate-normal saline (MS-NS), standard nest-oxytocin (SN-OXT), maternal separate-oxytocin (MS-OXT), maternal separate-cohabitation with healthy littermates (MS-COHAB)

Tail Suspension Test (TST)

Analysis of variance for the rate of immobility time showed no significant difference ($P = 0.84$). (Figure 4)




Figure 4. TST Rate of immobility time

Results from a 5 min TST. Rate of immobility time in 5 minutes.

Data are presented as means \pm SEM.

*: $p < 0.05$

Abbreviations: standard nest-normal saline (SN-NS), maternal separate-normal saline (MS-NS), standard nest-oxytocin (SN-OXT), maternal separate-oxytocin (MS-OXT), maternal separate-cohabitation with healthy littermates (MS-COHAB)

Social Interaction Test (SIT)

Significant differences were found in the analysis of variance for the rate of change in the low interaction zone stay ratio ($P < 0.01$). MS-NS showed a significant increase compared to SN-NS, MS-OXT and MS-COHAB ($P < 0.01$, $P = 0.0292$, $P < 0.01$). (Figure 5A)

SN-OXT showed no significant differences compared to SN-NS ($P = 0.9217$). (Figure 5B)




Figure 5. SIT Rate of change in the low interaction zone stay ratio

Results from SIT. Rate of change in the low interaction zone stay ratio with an unfamiliar mouse in the social interaction cage relative to that without the mouse. Data are presented as means \pm SEM.

*: $p < 0.05$

*: $p < 0.05$

Abbreviations: standard nest-normal saline (SN-NS), maternal separate-normal saline (MS-NS), maternal separate-oxytocin (MS-OXT), maternal separate-cohabitation with healthy littermates (MS-COHAB), standard nest-oxytocin (SN-OXT)

RT-PCR

In the comparison of *OXT* mRNA expression in the hypothalamus with *ACTB* as an endogenous control, a significant difference was found in the analysis of variance ($P < 0.01$). MS-NS, MS-OXT, MS-COHAB, and SN-OXT showed significant decreases compared with SN-NS ($P = 0.0284$, < 0.01 , 0.0144 , < 0.01). MS-OXT showed no significant difference compared with MS-NS ($P = 0.84$). MS-COHAB showed no significant difference compared

to MS-NS ($P = 0.10$). (Figure 6A)

In the comparison of *OXTR* mRNA expression in the hypothalamus with *ACTB* as an endogenous control, no significant difference was observed in the analysis of variance ($P = 0.87$). (Figure 6B)




Figure 6. Relative expression level of *OXT* and *OXTR* to *ACTB*

Results from RT-PCR. A: Relative expression level of *OXT* to *ACTB*, B: Relative expression level of *OXTR* to *ACTB*.

Data are presented as means \pm SEM.

*: $p < 0.05$

Abbreviations: standard nest-normal saline (SN-NS), maternal separate-normal saline (MS-NS), standard nest-oxytocin (SN-OXT), maternal separate-oxytocin (MS-OXT), maternal separate-cohabitation with healthy littermates (MS-COHAB)

Discussion

This study suggested that the administration of oxytocin and cohabitation with healthy littermates during adolescent to maternal-separated mice improved social behavior in the adult stage. The SIT showed a significant increase in MS-NS relative to SN-NS with respect to rate of change in the low interaction zone stay ratio. This indicates that maternal separation in the juvenile stage of the mice resulted in reduced social behavior in the adult stage, which is similar to earlier literature²⁰⁾. The SIT showed a significant difference between MS-NS and MS-OXT, and showed no significance between MS-OXT and SN-NS. This elucidated that oxytocin administration during adolescent improves adult socialization in maternal-separated mice. The SIT showed a significant difference between MS-NS and MS-COHAB, and showed no significance between MS-COHAB and SN-NS. This indicates that cohabitation with healthy mice improved social behavior.

The result of OFT showed significant differences between MS-NS and MS-COHAB, however, no conclusions could be drawn because there was no significant difference between SN-NS and MS-NS, or SN-NS and MS-COHAB. The OFT also showed significant differences between MS-OXT and MS-COHAB, however, it was not possible to conclude because there was no significant difference between SN-NS and MS-NS, and both groups were not significantly different from SN-NS. The results of the OFT showed movement is recognized in all individuals. We confirmed that no individuals were unable to move by the burden of maternal separation. In OFT, EPM, and TST, there was no significant difference between SN-NS and MS-NS. This result differs from those of previous studies on rodents¹³⁾. In order to increase the stress burden because of maternal separation, early weaning was also performed, similar to the highest stress burden according to previous literature¹²⁾. However, no significant differences in anxiety- or depression-associated behaviors were observed. This suggests the possibility that detailed rearing conditions not described in the literature, such as the time of experiment and the order of behavioral analysis, may affect the results in mice compared to rats, and the results were similar to those of previous systematic reviews with different results for rats and mice²¹⁾.

In our biological analysis, a comparison of mRNA expression of *OXT* in the hypothalamus with *ACBT* as

an intrinsic control showed a significant decrease in MS-NS compared to SN-NS, similar to results of existing literature²²⁾. Although the biological mechanism of the sociability changes induced by drug administration and environmental intervention was of interest, the biological analysis did not reveal significant changes in *OXT* and *OXT*R mRNA in the hypothalamus in either group, living with siblings and receiving oxytocin, and the mechanism could not be elucidated. There is literature indicating that there are sex differences in changes in the oxytocin system in the brain after maternal separation²³⁾. In this study, there were not enough individuals to separate the group by sex, and there is a possibility of not separated by sex affected the biological results. Further studies including larger-scale sex-specific experiments and biological comparisons in other brain regions, such as the AMY, PFC, Hip and NAc with reference to previous literature are needed^{24), 25)}.

MS-COHAB started cohabitation with healthy littermates after weaning not adolescent in order to avoid a decrease in affinity due to temporary separation with siblings. Therefore, it is possible that a significant difference was tend to occur compared to MS-OXT, in which intervention was performed only during adolescent. Several mice died because of the burden of maternal separation, which may have affected the results of behavioral analysis.

This study showed that oxytocin administration during adolescence and cohabitation with healthy littermates improves social behavior in maternal-separated mice. However, in the biological analysis, no significant differences in the expression of *OT* mRNA or *OTR* mRNA in the hypothalamus by each intervention.

Acknowledgments

We would like to thank Editage (www.editage.com) for English language editing.

References

- 1) Javakhishvili M, Widom CS: Childhood Maltreatment, Sleep Disturbances, and Anxiety and Depression: A Prospective Longitudinal Investigation. *Journal of Applied Developmental Psychology* 77: 101351, 2021.
- 2) Dube SR, Felitti VJ, Dong, M, Chapman DP, Giles WH, Anda RF: Childhood abuse, neglect, and

- household dysfunction and the risk of illicit drug use: the adverse childhood experiences study. *Pediatrics* 111: 564-572, 2003.
- 3) Porter C, Palmier-Claus J, Branitsky A, Mansell W, Warwick H, Varese F: Childhood adversity and borderline personality disorder: a meta-analysis. *Acta Psychiatrica Scandinavica* 141: 6-20, 2020.
 - 4) Crow TM, Levy KN, Bradley B, Fani N, Powers A: The roles of attachment and emotion dysregulation in the association between childhood maltreatment and PTSD in an inner-city sample. *Child Abuse Neglect* 118: 105139, 2021.
 - 5) Cecil CAM, Zhang Y, Nolte T: Childhood maltreatment and DNA methylation: A systematic review. *Neuroscience & Biobehavioral Reviews* 112: 392-409, 2020.
 - 6) Bosch OJ, Young LJ: Oxytocin and Social Relationships: From Attachment to Bond Disruption. *Current Topics in Behavioral Neurosciences* 35: 97-117, 2018.
 - 7) Bowen MT, Carson DS, Spiro A, Arnold JC, McGregor IS: Adolescent oxytocin exposure causes persistent reductions in anxiety and alcohol consumption and enhances sociability in rats. *PLoS One* 6: e27237, 2011.
 - 8) Tractenberg SG, Levandowski ML, Azeredo L A, Orso R, Roithmann LG, Hoffmann ES, Brenhouse H, Grassi-Oliveira R: An overview of maternal separation effects on behavioural outcomes in mice: Evidence from a four-stage methodological systematic review. *Neuroscience & Biobehavioral Reviews* 68: 48-503, 2016.
 - 9) Joushi S, Esmaeilpour K, Masoumi-Ardakani Y, Esmaeili-Mahani S, Sheibani V: Intranasal oxytocin administration facilitates the induction of long-term potentiation and promotes cognitive performance of maternally separated rats. *Psychoneuroendocrinology* 123: 105044, 2021.
 - 10) Tribollet E, Charpak S, Schmidt A, Dubois-Dauphin M, Dreifuss JJ: Appearance and transient expression of oxytocin receptors in fetal, infant and peripubertal rat brain studied by autoradiography and electrophysiology. *Journal of Neuroscience* 9: 1754-1773, 1989.
 - 11) Yamasaki A, Shinji T: Tips for Helping Children with Attachment Issues in School. *Journal of Learning Science Issue* 10: 49-55, 2017.
 - 12) Portero-Tresserra M, Gracia-Rubio I, Cantacorps L, Pozo OJ, Gomez-Gomez A, Pastor A, Lopez-Arnau R, de la Torre R, Valverde O: Maternal separation increases alcohol-drinking behavior and reduces endocannabinoid levels in the mouse striatum and prefrontal cortex. *European Neuropsychopharmacology* 28: 499-512, 2018.
 - 13) Kuniishi H, Ichisaka S: Early deprivation increases high-leaning behavior, a novel anxiety behavior, in the open field test in rats. *Neuroscience Research* 123: 27-35, 2017.
 - 14) Pellow S, Chopin P, File SE, Briley M: Validation of open-closed arm entries in an elevated plus-maze as a measure of anxiety in the rat. *Journal of Neuroscience Methods* 14: 149-167, 1985.
 - 15) Can A, Dao DT, Terrillion CE, Piantadosi SC, Bhat S, Gould TD: The tail suspension test. *Journal of Visualized Experiments* 59: e3769, 2012.
 - 16) Kaidanovich-Beilin O, Lipina T, Vukobradovic I, Roder J, Woodgett JR: Assessment of social interaction behaviors. *Journal of Visualized Experiments* 48: e2473, 2011.
 - 17) Feng T, An S, Kinden R, Zhang X, Jia R, Tai F: Alteration in oxytocin levels induced by early social environment affects maternal behavior and estrogen receptor alpha in mandarin voles (*Microtus mandarinus*). *Behavioural Brain Research* 365: 36-47, 2019.
 - 18) Lesse A, Rether K, Groger N, Braun K, Bock J: Chronic postnatal stress induces depressive-like behavior in male mice and programs second-hit stress-induced gene expression patterns of OxtR and AvpR1a in Adulthood. *Molecular Neurobiology* 54: 4813-4819, 2017.
 - 19) Sinclair MS, Perea-Martinez I, Dvoryanchikov G, Yoshida M, Nishimori K, Roper SD, Chaudhari N: Oxytocin signaling in mouse taste buds. *PLoS One* 5: e11980, 2010.
 - 20) Ryabushkina YA, Reshetnikov VV, Bondar NP: Maternal Separation Early in Life Alters the Expression of Genes Npas4 and Nr1d1 in Adult Female Mice: Correlation with Social Behavior. *Behavioural Neurology* 2020: 7830469, 2020.
 - 21) Wang, D, Levine JLS, Avila-Quintero V, Bloch M, Kaffman A: Systematic review and meta-analysis: effects of maternal separation on anxiety-like behavior in rodents. *Translational Psychiatry* 10: 174, 2020.
 - 22) Babygiriya R, Yoshimoto S, Gribovskaia-Rupp I, Bulbul M, Ludwig K, Takahashi T: Social interaction attenuates stress responses following chronic stress in maternally separated rats. *Brain Research* 1469: 54-62, 2012.
 - 23) Tanaka K, Osako Y, Takahashi K, Hidaka C, Tomita K, Yuri K: Effects of post-weaning social isolation on

- social behaviors and oxytocinergic activity in male and female rats. *Heliyon* 5: e01646, 2019.
- 24) Perkeybile AM, Carter CS, Wroblewski K L, Puglia MH, Kenkel WM, Lillard TS, Karaoli T, Gregory SG, Mohammadi N, Epstein L, Bales KL, Connelly JJ: Early nurture epigenetically tunes the oxytocin receptor. *Psychoneuroendocrinology* 99: 128-136, 2019.
- 25) Onaka T, Takayanagi Y: The oxytocin system and early-life experience-dependent plastic changes. *Journal of Neuroendocrinology* 33: e13049, 2021.
(Received on August 29th, 2022; Accepted on October 3rd, 2022)
[The authors declare no conflict of interest.]